

N<sup>o</sup> 1.

#14

Dr. James.

Peritonitis

Enteritis

Diarrhea

Erysipelas

Hematuria

On Erysipelas, Hysteritis

by Hardin Mays

Recd Mar 4<sup>th</sup> 1817.

On Cypripedium  
by H. B. K. & J. R. S.  
1847

A

Dissertation

in

Erysipelas

Capital

Capital

To show, and  
in fact,  
in the  
of some and  
a large proportion  
of the population.  
reference to  
to sympathy  
of the  
and of moral  
attached by  
indirectly,  
the cause  
is interesting  
product, and  
effects, that  
acted, in  
has done so  
has not been  
hence, as has  
and prejudicial

## Preliminary Remarks

The Skin, which has been denominated the parietes of the human body, considered physiologically, answers a fourfold purpose in the animal economy: It is the organ of touch; it covers and protects the whole structure; it is the outlet for a large proportion of the insensible perspiration, and it performs absorption. So extremely sensible is this part, such are its exposures to the attack of noxious agents, and so invariable are its sympathies with the whole body, and the various component organs, that we shall not be surprised, to find it often the seat of morbid affections. It is not infrequently the point just attacked by those noxious agents or causes, which either directly or indirectly, assail the life or health of man. So various are these causes of cutaneous diseases in their mode of operation; so intermingled and obscure are they in the effects which they produce, and such is the near resemblance very often of these effects, that no small degree of difficulty and perplexity is presented, in attempting to distinguish them. Anatomy, which has done so much in explaining many other morbid affections, has not been so successful in elucidating those of the skin; hence, as has been observed, the treatment of them is often empirical and prejudicial. Parts of an identity of structure are generally



affected by diseases very similar in appearance, but we are not from  
this circumstance, too hastily to conclude that they are the same;  
and to some more particularly will this remark apply, than those just  
mentioned. They are mostly eruptions of various kinds, possessing  
some characteristics in common, but differing in many others: thus  
a urticaria in small pox, encloses to each seems to possess a peculiar  
specific action producing effects sui generis. — On what this specific  
action depends I am unable to say; it is apparently a secretion of a  
particular kind which takes place according to some law of the animal  
economy, not yet explained. Without, however, our being able to com-  
prehend this subject, by attending carefully to the history of the various  
phenomena which occur in catarrhus diseases, we shall be enabled to  
treat them, in many cases successfully. Two circumstances seem  
instantly to influence them, and ought always to be kept in view  
viz. the structure of the part affected, and the nature of the various  
agent producing that effect.

Having premised these remarks, I proceed to the consideration  
of the subject, which is more particularly, the object of this essay.

...ing  
...ular  
...age, wh  
...ster  
...tion is  
...ejected  
...e genera  
...rally  
...temate  
...shall  
...elected  
...milar  
...esive  
...in com  
...ould as  
...stance,  
...e enjoy  
...e have  
...the sea  
...posed,  
...olutely

## Erysipelas.

According to a late writer, this may be defined "An inflammatory, cutaneous, and trivially elevated swelling, attended with redness, which disappears and leaves a white spot for a short time after being touched with the end of the finger; and the affection is characterized by a remarkable propensity, to spread with rapidity to a large extent." Although this disease has been generally considered an inflammatory affection, and it undoubtedly is so, in a very general acceptance of the term inflammation, yet on examining its symptoms more accurately we shall find it very different from phlegmon; the pain, the secretion, the whole of the phenomena, are in fact totally dissimilar. In erysipelas genuine pus is never secreted, adhesive inflammation does not circumscribe the disease as in common phlegmon, but matter once extravasated, travels sometimes to a great extent through the cellular substance, causing gangrene and trophication.

As erysipelas never appears in any other part but the skin we have a right to conclude, that this membrane alone is its seat; whether in the rete mucosum, as some have supposed, or the cutis vera, as stated by others, is not absolutely known.

order to ap-  
the history of  
writers have  
to be appea-  
sometimes:-  
list of Latin  
Hollan & a  
of the shire  
then the ap-  
of the exten-  
from cases  
sympelato-  
tion is an  
of the whole  
the most c  
the fongra  
these severa  
of the sam  
who have  
distinct c  
perhaps

43  
order to afford more precision, and greater accuracy, in  
the history of the symptoms and treatment of erysipelas.  
Writers have divided it into several species, according  
to the appearance of the local affection or constitutional  
symptoms:— of these the first and most general, is  
that of Idiopathic and Symptomatic.

Ballou says that when the disease is an affection  
of the skin alone, and very little of the whole system, "  
when the affection of the system is only symptomatic  
of the external inflammation, he calls it *Erythema*, a  
term used by Hippocrates to designate every kind of  
erysipelatous redness: but when the external inflamma-  
tion is an *eccanthema* and symptomatic of an affection  
of the whole system, *Erysipelas*. Besides this division  
the most common, are the *phlegmonous*, the *Edematous*,  
the *Gangrenous*, the *Eczatic*, and the *Bilious Erysipelas*.  
These several terms are expressive of the peculiar state  
of the same disease, and are not designed by those  
who have used them, to convey the idea of so many  
distinct complaints. The specific erysipelatous action  
presents always, its own peculiar characteristics.

At however  
nature  
of these for  
the most  
of the dis  
only as  
such as  
plaints,  
head acci  
from eve  
my good  
wonder m  
the eyes  
and oth  
this is  
and son  
with  
always  
When  
at m  
of the p

As however the affection requires some peculiarity of treatment, according as it assumes one or the other of these forms, it is proper to adopt them.

The most regular, as well as, the most ordinary form of the disease, and of which the others may be considered only as varieties, produced by accidental circumstances such as might have that effect with many other complaints, is the phlegmonous, affecting chiefly the head and face. As I conceive a description of this form will afford a better view of the disease, than any general account, I shall begin with it, and afterwards make some remarks on the others.

The erysipelas of the face comes on with a cold shivering, and other symptoms of pyrexia. The hot stage of this is frequently attended with a confusion of head and some degree of delirium, and almost always with ~~the~~ drowsings, and perhaps coma. The pulse is always frequent, and commonly full and hard. When these symptoms have continued, for one, two, or at most three days, there appears on some part of the face a redness: this at first is of no great extent

...gradual  
...the other  
...stated the  
...roads o  
...est of the  
...appears.  
...cribed.  
...at the  
...had con  
...the whole  
...plish a  
...the ex  
...posed to  
...later, be  
...parts of  
...almost c  
...in the re  
...in the be  
...his best  
...to discov  
...in the

6  
but gradually spreads from the part it first occupied  
to the other parts of the face, commonly till it has  
affected the whole; and frequently from the face it  
spreads over the hairy scalp, or descends on some  
part of the neck. As the redness spreads, it commonly  
disappears, or at least decreases in the parts, it has before  
occupied. All the parts on which the redness appears  
are at the same time affected, with some swelling,  
which continues for some time; after the redness has abated  
the whole face becomes considerably turgid, and the  
eyelids are often so much swelled, as entirely to shut  
up the eyes. When the redness and swelling, have  
proceeded for some time, there commonly arise sooner or  
later, blisters of a larger or smaller size, on several  
parts of the face. These contain a thin yellowish or  
almost colourless fluid, which is sooner, or later discharged  
by the rupture of the vesicles. The surface of the skin  
in the blistered places, sometimes becomes livid and black-  
ish, but this livor seldom goes deeper than the surface,  
or discovers any degree of gangrene affecting the skin.  
On the parts of the face not affected with blisters, the

...side se  
...quament  
...at  
...a  
...s  
...ing.  
...infla  
...for  
...it co  
...delic  
...increas  
...month  
...then the  
...time afte  
...the fac  
...has,  
...and with  
...to his or  
...the reso  
...the succ  
...covered

cuticle suffers towards the end of the disease, a considerable desquamation. Sometimes the tumor of the eyelids ends in suppuration. The inflammation coming upon the face does produce any remission of the fever which had before prevailed; and sometimes the fever increases, with the increasing, and spreading inflammation.

The inflammation usually continues for eight or ten days, and for the same time, the fever and symptoms attending it continue. In the progress of the inflammation, the delirium and coma attending it, sometimes go on increasing, and the patient dies apoplectic, on the seventh, ninth, or eleventh day of the disease.

When the fatal event does not take place, the inflammation after having affected a part, commonly the whole of the face, and perhaps the other external parts of the head, ceases. With the inflammation, the fever also ceases and without any evident crisis, the patient returns to his ordinary state of health.

The vesicles usually break on the fifth or sixth day, when the swelling subsides in some degree, leaving the surface covered with a yellowish, or black, irregular scab.

to pain wh  
as every other  
showing p  
water. In  
drop, while  
whiding  
perhaps, a  
then the  
was with  
day or two  
ation app  
from the p  
living re  
with a dis  
tain so  
best can  
day, large  
they soon  
and Lyon,  
of exten  
melled, a

The pain which attends the local affection of this, as well as every other form of erysipelas, is of a peculiar kind, a burning pain very much like that produced by boiling water. In some cases the disease terminates in a few days, while in others it continues much longer, or after subsiding in some degree, is again renewed with perhaps, aggravated symptoms.

When the disease affects the extremities, it usually commences with some febrile symptoms, which continue for a day or two, when some degree of redness and tumefaction appears on the limb; if on the leg, it extends from the foot to the knee. Its colour is an intense shining red, nearly continuous; it is also attended with a sensation of burning heat, and with excruciating pain, so that no pressure, not even the weight of a sheet can be borne upon it. On the third or fourth day, large vesications appear all over the part affected. They soon break and discharge a large quantity of serous lymph, and in some cases ulceration, more or less extensive, takes place. The leg is enormously swelled, and most painful during the night.

... some ...  
... on ...  
... some ...  
... to ...  
... it ...  
... very ...  
... matter ...  
... structure ...  
... the ...  
... I ...  
... the ...  
... extremities ...  
... species ...  
... shining ...  
... pebbles ...  
... free ...  
... St. ...  
... thick ...  
... chief ...  
... tilted ...  
... on ...

97

In some cases, the disease has been observed, after subsiding on one leg, to affect the other pretty much in the same manner, and in some cases it has been supposed to be translated to the brain, or other vital organs. When it affects the superior extremities, the symptoms are very similar to those just stated. In these cases matter is occasionally deposited, in the cellular structure.

The swelling of the glands, which is noticed by writers, I conceive to be a sympathetic affection.

The Erysipelas Edematous, affects the face, breast, or extremities. It has not so distinct an appearance as the species just mentioned: the swelling has a smooth shining surface, and is of a paler color, the pulse is feeble, and other signs of debility exist. The face resembles a bladder distended with water.

Mr Pearson, observes that, this form of the disease may attack persons of any age or temperament, but those are chiefly affected by it, whose constitutions are debilitated by age or excess; it is also frequently met with in dropical persons, in children and in new born

defects.  
cause or  
little to  
with a g  
conting  
Epistola  
of heat,  
of the s  
slight m  
has a r  
is of a  
with or  
constitut  
The Ego  
after Sec  
under the  
of  
many co  
Presides  
expies  
one part

Infants. — The Bilious Erysipelas of Doanet comes on most commonly, he states, with loss of appetite, bitter taste in the mouth, the tongue is covered with a yellow fur, nausea, and sometimes bilious vomitings supervene. The patient being debilitated, feels lassitude, wandering pains, and a sharp sensation of heat, without however, having a great dryness of the skin or much thirst. The tumefaction is slight most frequently insensible. The skin has a rose color bordering upon a yellow, the pain is of a smarting burning kind. It may commence with or without fever. This he considers as constituting properly the erysipelas of the Ancients. The Erysipelas affecting Infants at an early period after birth, was first described by Dr Underwood under the name of Anomalous inflammation of Infants.

The Zona or Shingles has been considered by many writers as a species of erysipelas. Besides these which are the more regular forms of erysipelas, it frequently occurs in patches, on various parts of the body, attended by little, or no fever.

if any found  
be this:  
from some  
most likely  
is another  
state of fe  
putrid, or  
the parts  
andification  
two are a  
may take  
fence. The  
take place  
me. in a  
Am  
specimens for  
satisfactory  
of Hippoc  
ed upon  
Hoson A  
for Violent

If any form of the disease be purely local, it must be this: under this might be included that arising from wounds.

Most authors mention the gangrenous state of erysipelas as another species. It seems to be attended by that state of febrile action, denominated by the old writers putrid, or what may now be Typhus gravior.

The parts exhibit the usual signs of gangrene and mortification, and frequently sloughs, resembling tooth-  
 town are discharged; but this sloughing I conceive, may take place without so malignant a state of fever. The serous collections of matter which often take place in the common acute form of the disease, in the cellular structure, may cause them.

Among the many causes which have been assigned for erysipelas, none seem very distinct or satisfactory. The most ancient opinion is that of Hippocrates and Galen who supposed it depended upon a congestion of the bile.

Modern authors enumerate the following,

1<sup>st</sup> Violent Passions. — 2 Exposure to the heat of the

due, or  
improper  
negotiable  
intention  
operation  
There can  
possibly  
see it  
there is  
mally s  
Is any  
on this s  
whether  
some imp  
without  
be person  
the law  
achieve a  
they gen  
attacks  
will its

sun, or that of fire too long continued, — 3<sup>d</sup> The  
impression of cold damp air, — 4<sup>th</sup> The action of various  
vegetable, animal, or mineral poisons, 5<sup>th</sup> Wounds  
contusions, fractures &c. 6<sup>th</sup> Cold feet, — 7<sup>th</sup> Suppressed  
evacuations. — 8<sup>th</sup> Disorder of the primæ viæ.

There can be no doubt that, erysipelas has sometimes  
prevailed as an epidemic: Dr. Parr says he has  
seen it four times; and Mr. Pearson observes that  
there is some reason to conclude that it is occasi-  
onally so. —

Is erysipelas contagious? The best Authorities  
on this subject seem not to decide positively,  
whether it is, or is not so. It certainly is of  
some importance to decide this question correctly,  
without however taking this upon myself, I may  
be permitted to remark, that it does not follow  
the laws of those diseases, which are known and  
acknowledged by every one, to be strictly contagious.  
They generally destroy the susceptibility to a second  
attack, whereas it is increased by this.  
Will its becoming Tophus render it contagious?

ought it o  
not crys  
moving f  
of conlage  
of this dis  
kind, is a  
which atten  
tutions of  
gods of the  
allow. It a  
by the loca  
said, ene  
may be, co  
mas char  
change fr  
his influen  
the type of  
age observ  
this compe  
By the com  
who suppo

Ought it not to produce typhus fever than, and not erysipelas? There can be no doubt of its occurring frequently when there is no possible chance of contagion. What then is the peculiar nature of this disease? Local inflammation of every kind, is a very distinct thing from the fever which attends it, and while we admit many distinctions of the former, the same will not hold good of the latter. At the same time we must allow, that the fever is often much influenced by the local disease. — From all that has been said, we may infer, that the febrile affection may be either, inflammatory, typhus, or of a mixed character, that it may, and often does change from one to the other. These forms may be influenced by several circumstances, such as the type of the prevailing epidemic; it was long ago observed by Sydenham, that like a desert at this compelled every disease to wear its livery. By the constitution and habits of the person, — those who support life, or rather force it, almost

actively  
material  
the loca  
Mineral  
prevails.  
The form  
nature, a  
from  
of this de  
respected  
highly imp  
very often  
opposite c  
happens of  
With  
a little disc  
since  
principally  
there have  
on other me  
present time

entirely by the Acute Stimulus, will of course very materially alter it. — The Season ~~of the~~ of the year — the local situation — the prevalence of Marsh Miasmata. In the fall, and where Malignant Fever prevails, it will assume much of its character. This form is mentioned particularly by the French writers, and by practitioners of warm climates.

From every view that I have been able to take of this disease, and from the opinions of the most respectable authority, I conclude, that it is of a highly inflammatory character, disposed to run, very often, if not checked, in a short time into an opposite extreme, and that the greatest mischief happens from an excess of action.

With regard to the treatment of erysipelas, not a little discordance has prevailed amongst practitioners ever since the days of Hippocrates, some depending principally on mesotocion, some on purgatives, while others have declared them both pernicious, and relied on other means. The European physicians, of the present time, are not at all agreed as to the best

work of the  
may be  
philology  
partitions  
to tonic  
acute  
quilt de  
forms of  
reason of  
one over  
well as  
circulation  
in tonic  
acute  
figure of  
of music  
Gustav  
persuade  
in every  
according  
not by

mode of treating erysipelas. The Dutch writers, among whom may be mentioned, M<sup>r</sup> Bell and V<sup>an</sup> Cullen, recommend the Antiphlogistic plan. M<sup>r</sup> Pearson and most of the London practitioners, use evacuations sparingly and resort early to tonics and stimulants, while the French adopt very moderate means to evacuate the bile, and produce a gentle diaphoresis. Now by attending to the various forms of the disease, which we have described, the reason of this difference in practice, will appear at once evident. In our own country, erysipelas as well as every other acute disease, requires copious evacuations, and while these are proper, all stimulants or tonics, must be manifestly improper. We not only evacuate more, but endeavor to produce the same degree of impression, have to use much larger doses of medicine, for however thoroughly, one grain of Emetic Tartar, might evacuate a Frenchman, we are persuaded, that it would require more for an American. In every case it is necessary, to treat the disease according to the existing state of the system, and not by the name.

The case  
in which  
is the  
situation, &  
of the d  
Employe  
have rec  
not only  
when rec  
route fo  
in favor  
of the m  
may me  
sufficient  
be from  
we have  
calculated  
one of M  
stimulus  
the disease  
objections

The curative means are generally divided into the general or constitutional, and the local.

As the most powerful means of reducing arterial action, and thereby moderating, or overcoming the violence of the disease, bloodletting may be placed first.

Employed indiscriminately, as Celsus and some others have recommended, it could not fail, in many cases, not only to be useful, but highly pernicious; but when resorted to at an early period, and in the acute form of the disease, it is an invaluable remedy. In favor of this practice, we might enumerate many of the most respectable practitioners, among whom we may mention Sydenham, and Rush, names amply sufficient, where authority is required.

In favor of Emetics, there is not so much authority; we have however, amongst several others, Richter, the celebrated professor of Gottingen, who is justly considered one of the most experienced, and judicious practitioners. It has been remarked by Parry, that the disease often terminates, by vomiting and bilious dejections, from which we may correctly infer that

on such  
lands or  
improvements  
system  
I cannot  
their acc  
Negatives  
they have  
and mean  
It is as  
to keep  
the phleg  
hanging  
hanging  
viz the  
cases be  
Various  
Hose co  
the Ant  
one gra  
this to

17  
in such cases, emetics would prove highly beneficial: besides evacuating the stomach, they produce a powerful impression, through its sympathies, on the whole system, and particularly on the skin: in these cases I cannot perceive any mischief likely to result from their use.

Purgatives have been highly extolled by some, while they have been as much opposed by others: - Galen and many since his time have recommended them.

It is allowed by every one to be proper in all cases to keep the bowels in a soluble state, and in the phlegmonous erysipelas, of the face and head, active purging may be placed next to venisection, but as purging interferes with another important indication viz the determination to the surface, it in many cases becomes a questionable remedy.

Various diaphoretics have been recommended, but those commonly preferred at the present time, are the Antimonials. Depault's practice was to give one grain of the Emetic Tartar, largely diluted, and this to be repeated if necessary.

Alst. Plumb.

In every  
nature, a  
to the  
up and  
the loca  
the gene  
have bee  
insuffice  
a little  
the flue  
however  
place a  
the use  
were us  
they ex  
bearing  
fell over  
they ex  
to that  
sped by

In every case, the body must be kept at a moderate temperature, such as will be most agreeable to the patient's feelings. In the Edematous and gangrenous erysipelas, we evacuate, and resort earliest to stimulants.

The local means have been, even more numerous than the general, various ointments, cataplasms, washes &c have been recommended, but have all been found inefficacious or prejudicial. Dr Bullen persuaded of this, that he discards them all, except a little flour sprinkled on the part, in order to absorb the fluid discharged from the vesicles. Since his time however, a very important improvement, has taken place in the treatment of this disease. I allude to the use of blisters, and a solution of opium. Blisters were used so long ago as the time of Ambrose Paré: they were then directed to be applied to the neighbouring parts. A very different practice is now followed; we apply them directly to the diseased part. They excite a new action in the part, and according to that law of the Animal Economy, so ably developed by M Hunter, the incompatibility of two actions of

equal for  
income  
healthy a  
abstraction  
out of exp  
readful  
tempting  
is it wou  
the face  
clusion of  
be the a  
is breise  
they shoul  
when a  
be made,  
jensen lo

equal force in the same part, at the same time.  
overcome the disease, and allow the part to assume its  
healthy action. To Dr. Physick we are indebted for the  
introduction of this excellent practice, in the treat-  
ment of erysipelas. Cases which formerly spread with  
dreadful devastation, can now be checked with as much  
promptness, and certainty, as any other disease.

As it would not be so proper to apply blisters to  
the face, it is in this case that we resort to the  
solution of opium, with great advantage.

In the erysipelas which so often attends wounds  
and bruises, blisters are eminently serviceable;  
they should be applied all over the diseased part.

When collections of matter form, free incisions should  
be made, and not small openings as some have advised,  
in order to evacuate the matter, with the sloughs

